

COMMUTATIVE ALGEBRA

YOSHIFUMI TSUCHIMOTO

02. Localization

DEFINITION 2.1. Let A be a commutative ring. Let S be its subset. We say that S is multiplicative if

- (1) $1 \in S$
- (2) $x, y \in S \implies xy \in S$

holds.

DEFINITION 2.2. Let S be a multiplicative subset of a commutative ring A . Then we define $A[S^{-1}]$ as

$$A[\{X_s; s \in S\}] / (\{sX_s - 1; s \in S\})$$

where in the above notation X_s is a indeterminate prepared for each element $s \in S$.) We denote by ι_S a canonical map $A \rightarrow A[S^{-1}]$.

LEMMA 2.3. Let S be a multiplicative subset of a commutative ring A . Then the ring $B = A[S^{-1}]$ is characterized by the following property:

Let C be a ring, $\varphi : A \rightarrow C$ be a ring homomorphism such that $\varphi(s)$ is invertible in C for any $s \in S$. Then there exists a unique ring homomorphism $\psi = \phi[S^{-1}] : B \rightarrow C$ such that

$$\varphi = \psi \circ \iota_S$$

holds.

COROLLARY 2.4. Let S be a multiplicative subset of a commutative ring A . Let I be an ideal of A given by

$$I = \{x \in I; \exists s \in S \text{ such that } sx = 0\}$$

Then I is an ideal of A . Let us put $\bar{A} = A/I$, $\pi : A \rightarrow \bar{A}$ the canonical projection. Then:

- (1) $\bar{S} = \pi(S)$ is multiplicatively closed.
- (2) We have

$$A[S^{-1}] \cong \bar{A}[\bar{S}^{-1}]$$

- (3) $\iota_{\bar{S}} : \bar{A} \rightarrow \bar{A}[\bar{S}^{-1}]$ is injective.

There is another description of $A[S^{-1}]$. Namely, We consider an equivalence relation \sim_S on a set $S \times A$ by

$$(s_1, a_1) \sim_S (s_2, a_2) \iff t(s_1a_2 - s_2a_1) = 0 (\exists t \in S)$$

We call the quotient space $S \times A / \sim_S$ as $S^{-1}A$. The equivalence class of $(s, a) \in S \times A$ in $S^{-1}A$ is denoted by $s^{-1}a$. Then it is easy to introduce a ring structure of $S^{-1}A$ and see that $S^{-1}A$ actually satisfies the universal property of $A[S^{-1}]$. We thus have a canonical isomorphism $S^{-1}A \cong A[S^{-1}]$.

EXAMPLE 2.5. $A_f = A[S^{-1}]$ for $S = \{1, f, f^2, f^3, f^4, \dots\}$. The total ring of quotients $Q(A)$ is defined as $A[S^{-1}]$ for

$$S = \{x \in A; x \text{ is not a zero divisor of } A\}.$$

When A is an integral domain, then $Q(A)$ is the field of quotients of A .

DEFINITION 2.6. Let A be a commutative ring. Let \mathfrak{p} be its prime ideal. Then we define the localization of A with respect to \mathfrak{p} by

$$A_{\mathfrak{p}} = A[(A \setminus \mathfrak{p})^{-1}]$$

DEFINITION 2.7. Let S be a multiplicative subset of a commutative ring A . Let M be an A -module we may define $S^{-1}M$ as

$$\{(m/s); m \in M, s \in S\} / \sim$$

where the equivalence relation \sim is defined by

$$(m_1/s_1) \sim (m_2/s_2) \iff t(m_1s_2 - m_2s_1) = 0 \quad (\exists t \in S).$$

We may introduce a $S^{-1}A$ -module structure on $S^{-1}M$ in an obvious manner.

$S^{-1}M$ thus constructed satisfies an universality condition which the reader may easily guess.

By a universality argument, we may easily see the following proposition.

PROPOSITION 2.8. *Let A be a commutative ring. Let S be a multiplicative subset of A . Let M be an A -module. Then we have an isomorphism*

$$S^{-1}A \otimes_A M \cong S^{-1}M$$

of $S^{-1}A$ -modules.

PROPOSITION 2.9. *Let A be a commutative ring. Let S be a multiplicative subset of A . Then the natural homomorphism $A \rightarrow S^{-1}A$ is flat.*

2.1. local rings.

DEFINITION 2.10. A commutative ring A is said to be a local ring if it has only one maximal ideal.

EXAMPLE 2.11. We give examples of local rings here.

- Any field is a local ring.
- For any commutative ring A and for any prime ideal $\mathfrak{p} \in \text{Spec}(A)$, the localization $A_{\mathfrak{p}}$ is a local ring with the maximal ideal $\mathfrak{p}A_{\mathfrak{p}}$.

DEFINITION 2.12. Let A, B be local rings with maximal ideals $\mathfrak{m}_A, \mathfrak{m}_B$ respectively. A local homomorphism $\varphi : A \rightarrow B$ is a homomorphism which preserves maximal ideals. That means, a homomorphism φ is said to be local if

$$\varphi^{-1}(\mathfrak{m}_B) = \mathfrak{m}_A$$

EXAMPLE 2.13 (of NOT being a local homomorphism).

$$\mathbb{Z}_{(p)} \rightarrow \mathbb{Q}$$

is not a local homomorphism.

.....

LEMMA 2.14 (Zorn's lemma). *Let \mathcal{S} be a partially ordered set. Assume that every totally ordered subset of \mathcal{S} has an upper bound in \mathcal{S} . Then \mathcal{S} has at least one maximal element.*